## **Heating Matter**

#### **Ouestions**

Q1.

A student uses the apparatus in Figure 17 to determine the specific heat capacity of water.




Figure 17

i) State the measurements needed to calculate the specific heat capacity of water.			
(4			

(ii) State	two ways that the apparatus could be adapted to improve the procedure.	
	(2	2)
1		
2		
	(Total for question = 6 marks	)

Q2.

A student wants to determine the specific heat capacity of copper.

Figure 20 shows a piece of copper, with a thread tied around it, in a glass beaker of boiling water.




Figure 20

The student leaves the piece of copper in the boiling water so that the copper reaches a temperature of 100 °C.

The student uses the thread to take the piece of copper out of the boiling water.

The student puts the hot piece of copper into a different beaker of cold water at 20 °C.

The apparatus is shown in Figure 21.




Figure 21

The student assumes that the thermal energy gained by the water equals the thermal energy lost by the piece of copper.

The water and copper both reach a temperature of 22 °C.

The cold water gains 1050 J of energy.

The mass of the piece of copper is 0.058 kg.

(i) Calculate a value for the specific heat capacity of copper, using these results.	
Use the equation	
change in thermal energy = mass × specific heat capacity × change in temperature	
$\Delta Q = m \times c \times \Delta \theta$	
(2)	)
specific heat capacity of copper from these results =	
J/kg °C	;
(ii) The value for the specific heat capacity of copper obtained from the student's results is	
lower than the correct value.	
State two ways that the experiment could be improved to give a value that is closer to the	۱e
correct value. (2)	1
1	
2	
(Total for question = 4 marks)	

A beaker contains 0.25 kg of water at room temperature. The beaker of water is heated until the water reaches boiling point (100 °C). The specific heat capacity of water is 4200 J/kg °C. The total amount of thermal energy supplied to the water is 84 000 J.	
(i) Calculate the temperature of the water before it was heated.	
Use an equation selected from the list of equations at the end of this paper	. (3)
town and the bafana basting	
temperature before heating =	°C
(ii) The heating continues until 0.15 kg of the water has turned into steam.	°C
(ii) The heating continues until 0.15 kg of the water has turned into steam.  The thermal energy needed to turn the boiling water into steam is 0.34 MJ.	
(ii) The heating continues until 0.15 kg of the water has turned into steam.	
(ii) The heating continues until 0.15 kg of the water has turned into steam.  The thermal energy needed to turn the boiling water into steam is 0.34 MJ. Calculate the specific latent heat of vapourisation of water.	:
(ii) The heating continues until 0.15 kg of the water has turned into steam.  The thermal energy needed to turn the boiling water into steam is 0.34 MJ. Calculate the specific latent heat of vapourisation of water.	:
(ii) The heating continues until 0.15 kg of the water has turned into steam.  The thermal energy needed to turn the boiling water into steam is 0.34 MJ. Calculate the specific latent heat of vapourisation of water.	:
(ii) The heating continues until 0.15 kg of the water has turned into steam.  The thermal energy needed to turn the boiling water into steam is 0.34 MJ. Calculate the specific latent heat of vapourisation of water.	. (2)

(iii) The graph in Figure 13 shows how the  $\,$  volume of 1 kg of water changes with temperature.




Figure 13

Describe how the density of water changes with temperature over the range of temperature shown in Figure 13. Calculations are not required.

(2)

(Total for question = 7 marks)

Q4.

(i) Figure 11 shows an electric kettle.




Figure 11

The kettle contains 1.5 kg of water.

The kettle is switched on.

Calculate the energy needed to raise the temperature of the water by 50 °C.

Specific heat capacity of water = 4200 J/kg °C

Use the equation

$$\Delta Q = m \times c \times \Delta \theta$$

(2)

(ii) The amount of energy, E, needed to bring the water to boiling point is 670 000 J.

The kettle has a power of 3500 W.

Calculate the time, t, it takes to bring the water to boiling point.

Use the equation

$$P = \frac{E}{t}$$

(3)

A digital thermometer gives a temperature reading of 23 °C.  Calculate the value of this temperature in kelvin.  (Total for question = 1 mark)  Q6.  An electric kettle contains 1.41 kg of water at 25 °C.  The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	Q5.	
Q6.  An electric kettle contains 1.41 kg of water at 25 °C. The kettle is switched on.  After a while, the water reaches boiling point at 100 °C. The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle. Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (ii) The kettle is kept switched on and the water continues to boil.  After a while, the mass of the water in the kettle has decreased to 1.21 kg. The thermal energy supplied to the water during this time was 450 000 J. Calculate the specific latent heat of vaporisation of water.  Use an equation selected from the list of equations at the end of the paper.	A digital thermometer gives a temperature reading of 23 °C.	
Q6.  An electric kettle contains 1.41 kg of water at 25 °C.  The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	Calculate the value of this temperature in kelvin.	
Q6.  An electric kettle contains 1.41 kg of water at 25 °C.  The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =		(1)
Q6.  An electric kettle contains 1.41 kg of water at 25 °C.  The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =		•••••
Q6.  An electric kettle contains 1.41 kg of water at 25 °C.  The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =		•••••
Q6.  An electric kettle contains 1.41 kg of water at 25 °C.  The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	(Total for question = 1	1 mark)
An electric kettle contains 1.41 kg of water at 25 °C.  The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =		·
The kettle is switched on.  After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	Q6.	
After a while, the water reaches boiling point at 100 °C.  The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle. Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	An electric kettle contains 1.41 kg of water at 25 °C.	
The specific heat capacity of water is 4200 J / kg °C.  (i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	The kettle is switched on.	
(i) Calculate the amount of thermal energy supplied to the water by the kettle.  Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	After a while, the water reaches boiling point at 100 °C.	
Give your answer to the appropriate number of significant figures.  Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =	The specific heat capacity of water is 4200 J / kg °C.	
Use an equation selected from the list of equations at the end of the paper.  (3)  energy supplied =		
energy supplied =		
(ii) The kettle is kept switched on and the water continues to boil.  After a while, the mass of the water in the kettle has decreased to 1.21 kg. The thermal energy supplied to the water during this time was 450 000 J. Calculate the specific latent heat of vaporisation of water.  Use an equation selected from the list of equations at the end of the paper.		(3)
(ii) The kettle is kept switched on and the water continues to boil.  After a while, the mass of the water in the kettle has decreased to 1.21 kg. The thermal energy supplied to the water during this time was 450 000 J. Calculate the specific latent heat of vaporisation of water.  Use an equation selected from the list of equations at the end of the paper.		
(ii) The kettle is kept switched on and the water continues to boil.  After a while, the mass of the water in the kettle has decreased to 1.21 kg. The thermal energy supplied to the water during this time was 450 000 J. Calculate the specific latent heat of vaporisation of water.  Use an equation selected from the list of equations at the end of the paper.		
(ii) The kettle is kept switched on and the water continues to boil.  After a while, the mass of the water in the kettle has decreased to 1.21 kg. The thermal energy supplied to the water during this time was 450 000 J. Calculate the specific latent heat of vaporisation of water.  Use an equation selected from the list of equations at the end of the paper.		7
After a while, the mass of the water in the kettle has decreased to 1.21 kg. The thermal energy supplied to the water during this time was 450 000 J. Calculate the specific latent heat of vaporisation of water. Use an equation selected from the list of equations at the end of the paper.		J
The thermal energy supplied to the water during this time was 450 000 J. Calculate the specific latent heat of vaporisation of water. Use an equation selected from the list of equations at the end of the paper.		
Use an equation selected from the list of equations at the end of the paper.  (3)	The thermal energy supplied to the water during this time was 450 000 J.	
specific latent heat of vaporisation =		(3)
specific latent heat of vaporisation =		
specific latent heat of vaporisation =		
- σρουπο ιαιοπι ποαι οι ναρυποαιιοπ J / κχ	specific latent heat of vanorisation =	1 / ka
(Total for question = 6 marks)		_

Q7.	
Another student decides to melt some ice.	
The student melts 380 g of ice at 0 °C.	
The specific latent heat of fusion of ice is 3.34 × 105 J/kg.	
Calculate the thermal energy needed to melt the ice.	
Select an equation from the list of equations at the end of this paper.	
	(2)
thermal energy needed =	7
thermat energy needed =	J
(Total for question	= 2 marks)
Q8.	
On a very cold day, the temperature of the air is -4 °C.	
Calculate the value of this temperature on the kelvin scale.	
	(1)
temperature =	K
(Total for question	า = 1 mark)

$\overline{}$	$\sim$		
. )	ч		
$\smile$	J	٠	

Describe, in terms of particles,	two differences	between a s	olid and a	a liquid of tl	ne same
substance.					

1			 (∠)
Δ	•••••	•••••	 •
	•••••	•••••	
2			
	•••••	•••••	

(Total for question = 2 marks)

Q10.

\* This question is about determining the specific heat capacity of aluminium. An aluminium block is placed in boiling water as shown in Figure 21.




Figure 21

The piece of string is tied to the aluminium block so the block can be transferred from the boiling water to the cold water.

Describe how a student could use this apparatus, and any additional items needed, to determine the specific heat capacity of aluminium.

Your answer should include how the student would

obtain the necessary measurements
use the measurements to calculate the specific heat capacity of aluminium.

(6)

(Total for question = 6 marks)

Q11.

The student decides to measure the temperature of the water every minute while it is being heated.

Figure 18 shows a graph of the student's results.




Figure 18

Predict the temperature of the water if the heating continues up to 8 minutes.

(1)

temperature of the water = .....°C

(Total for question = 1 mark)

Q12.

Figure 21 shows a shed made mostly of concrete blocks.




Figure 21

State two practical ways to reduce heat loss from this shed.

4				(∠)
⊥	 •••••	•••••	•••••	•••••
2	 			

(Total for question = 2 marks)

# Mark Scheme – Heating Matter

Q1.

Question number:	Answer Additional guidance		Mark
(i)			(4) AO 1 2
	(measurement of) the mass of water (1)	accept volume / weight of water ignore amount	
	(measurement of) the temperature (rise/change) (1)	accept (take) thermometer reading	
	(measurement of) the energy supplied / from heater (1)	accept (take) reading of the joulemeter ignore 'change in thermal energy' (from equation)	
	detail of any of the above (1)	e.g. measure temp at the start and end or measure mass of empty cup or start and end readings on the meter	

Question Number:	Answer	Additional guidance	Mark
(ii)	any two improvements from:	both marks can be scored in one answer space	(2) AO 3 3b
		ignore repeating readings ignore increase voltage / power / energy ignore use of clamp to hold thermometer / heater	
	add lid /cover (1)		
	add lagging / insulation (1)	accept use better insulator or better insulated / thicker cup accept use calorimeter	
		ignore use glass beaker unless cup is inside it ignore different type of cup	
	add a stirrer (1)		
	use a more sensitive thermometer (1)	accept use digital / electric thermometer / data logger	
	ensure heater fully submerged (1)		

#### Q2.

Question number	Answer	Additional guidance	Mark
(i)	rearrangement (and substitution) (1)		(2)
	(c) = $\frac{1050}{0.058 \times 78}$	$c = \Delta Q$ $m \times \Delta \theta$	
		award 1 mark if 78 seen	
	evaluation (1) 230 (J/kg °C)	accept 232(J/kg °C)	
		award full marks for correct answer without working.	

Question number	Answer	Additional guidance	Mark
(ii)	any two of the following	ignore more accurate measurements e.g. thermometer, balance etc. ignore taking repeats	(2)
	reduce heat loss from water/insulate beaker/add cover (1)		
	make the temperature rise larger/use a larger piece of copper/ use a smaller amount of water (1)	start with colder water	
	(use)a stirrer (1)		
	account for heat gained by glass beaker (1)		
	transfer the hot copper faster (1)		
	use a different heating method (1)		
	measure the temperature of the boiling water (1)		

Q3.

Question Number	Answer		Additional guidance	Mark
(i)	substitution into $\Delta Q = m \times c \times \Delta$	9	accept	(3)
	(1)		substitution and	- 1
	84 000 = 0.25 × 4200 × Δθ		rearrangement in either order	
	0.25 4200 20		in either order	
	rearrangement $\frac{\Delta Q}{m \times c}$ (1)			
	$(\Delta \theta = ) \frac{84000}{0.25 \times 4200}$			
	(= 80)			
	evaluation (1)			
	(temperature before heating = 20 (°C)	)	answer of 80 (°C) scores 2 marks	
	20 ( C)			
			award full marks	
			for the correct	
			answer without	
Overtion	0	۸ ما ما	working	Mayle
Question Number	Answer	Add	itional guidance	Mark
(ii)	substitution into $Q = m \times L(1)$			(2)
	0.34 = 0.15 × L			
	re-arrangement and evaluation (1)			
	$(L = \frac{0.34}{0.15} = )$			
	2.3 (MJ/kg)		w values that nd to 2.3 (MJ/kg)	
		allov erro	w 1 mark for POT r	
		the	rd full marks for correct answer out working	

Question Number	Answer	Additional guidance	Mark
(iii)	A description that makes reference to any <b>two</b> of the following		(2)
	(density) increases between 0°C and 4°C (1)	increases initially / at first / up to 4°C	
	reaches a maximum at 4°C (1)		
	(density) decreases above 4 °C (1)	then decreases	
		if no other marks scored then credit reference to large volume means low density (OWTTE) for 1 mark only	

#### Q4.

Question	Answer	Additional guidance	Mark
Number			
(i)	substitution (1)		(2)
	(ΔQ) = 1.5 x 4200 x 50		
		0.50	
	evaluation (1)	accept 315 000 (J)	
	320 000 (J)	310 000 (J)	
		award full marks for the	
		correct answer without	
		working	
		320 000 000	
		315 000 000	
		310 000 000 score 1 mark	
		(mass in grams)	

Question	Answer	Additional guidance	Mark
Number			
(ii)	substitution (1)	accept substitution and	(3)
	3500 = <u>670 000</u>	rearrangement in either	
	t	order	
	rearrangement (1)		
	(t=) <u>670 000</u>		
	3500		
	evaluation (1)		
	190(s)	accept any answer that	
		round to 190(s)	
		power of ten error award 2	
		marks maximum	
		award full marks for the	
		correct answer without	
		working	

#### Q5.

Question	Answer	Additional	Mark
Number		guidance	
	296 (°C)	accept 23 + 273	(1)

#### Q6.

Question number	Answer	Additional guidance	Mark
i	substitution into $\Delta Q = m \times s \times \Delta T \qquad (1)$		(3) AO2
	$(\Delta Q) = 1.41 \times 4200 \times (100-25)$		

	1 3 0 3,703	2 8 38	ignore POT error for this mark
6	evaluation	(1)	
	(energy =) 444,15	50 (J)	
a	answer to 2 sf	(1)	
4	440,000 (J)		independent mark allow 3 sf 444,000
			award full marks for the correct answer without working
			award 1 mark for answers with values 148,050 or 592,200 (incorrect temp and sf)
			award 2 marks for answers with values 150,000 or 148,000 or 590,000 or 592,000 (incorrect temp but allowed sf)

Question number	Answer		Additional guidance	Mark
ii	substitution into $\Delta Q = m \times L$ $450,000 = (1.41 - 1.21) \times L$ rearrangment	1)	allow substitution and rearrangement in either order	(3) AO2
	evaluation (L) = 2 200 000 (J/kg)	1)	accept 2 250 000  award full marks for the correct answer without working  award 1 mark for answers that round to 330,000 or 370,000 (incorrect mass used)	

#### Q7.

Question Number:	Answer	Additional guidance	Mark
	substitution (1) $(Q =) \frac{380 \times 3.34 (\times 10^5)}{(1000)}$		(2) AO 2 1
	evaluation (1)		
	1.27 x 10 <sup>5</sup> (J)	127 kJ 126920 (J)	
		accept answers that round to 1.27 x 10 <sup>5</sup> e.g. 1.2692 x 10 <sup>5</sup>	
		accept 130 kJ or 1.3 x 10 <sup>5</sup> (J)	
		POT error max. 1 mark	
		award full marks for correct answer without working	

#### Q8.

Question number	Answer	Additional guidance	Mark
	269 (K)	allow use of 273.14? 269.14 (K)	(1) AO2

#### Q9.

Question number	Answer	Additional guidance	Mark
	descriptions to include any <b>two</b> of		(2) AO1
	particles / atoms in solid close(r) together (1)	reverse argument	
		difference asked for, so must compare for subsequent marking points	
	particles / atoms in solid (vibrate) in fixed positions but particles in liquid move (freely) (1)		
	particles in a solid in regular arrangement but particles in liquid are randomly arranged (1)		
	particles in a liquid have more (kinetic) energy (than in a solid) (1)		
		allow answers in terms of forces between particles	

## Q10.

Question number	Indicative content	Mark
*	Answers will be credited according to candidate's deployment of knowledge and understanding of the material in relation to the qualities and skills outlined in the generic mark scheme.  The indicative content below is not prescriptive and candidates are not required to include all the material which is indicated as relevant. Additional content included in the response must be scientific and relevant.  Procedure  • Measure the temperature of the boiling water  • Allow sufficient time for block to reach temperature of boiling water  • Measure temperature of cold water in beaker  • Using a thermometer  • Transfer (hot) aluminium block to cold water in the beaker.  • Work quickly to avoid thermal energy loss during transfer  • Measure temperature of water  • Stir to ensure even distribution  • Measure maximum temperature reached by water  • Calculate temp rise of water by subtracting initial from final temperature.  • Calculate temp drop of aluminium by subtracting final temperature from 100.  • Find mass of beaker and water and aluminium  • Use a balance  • Empty water from beaker and dry beaker and block  • Weigh beaker and block alone  • Find mass of water by subtraction.  • Allow plausible method of finding mass of water before putting block in.	(6) AO2 and AO3
	Process results  • Calculate thermal energy gained water using ΔQ = m x c × Δθ  • Thermal energy gained by water = thermal energy lost by aluminium  • Specific heat capacity of aluminium =   thermal energy transferred mass of Al × temp drop of Al	

Level	Mark	Descriptor		
	0	No awardable co	ontent	
Level 1	1–2	understanding of flawed or simplicontext of the of Analyses the so-connections are		
Level 2	3-4	The plan is mos knowledge and and procedures elements in the  Analyses the sc connections bet procedures. A p	tly supported through linkage and application of understanding of scientific enquiry, techniques, some logical connections made between context of the question. (AO2) ientific information and provides some logical ween scientific enquiry, techniques and artially completed plan that synthesises mostly tanding, but not entirely coherently. (AO3)	
Level 3	5-6	knowledge and and procedures the context of the context of the seconnections between the connections and the connections are t	The plan is supported throughout by linkage and application of knowledge and understanding of scientific enquiry, techniques and procedures, logical connections made between elements in the context of the question. (AO2)  Analyses the scientific information and provide logical connections between scientific concepts throughout. A well-developed plan that synthesises relevant understanding	
Level	Mark	Additional Guidance	General additional guidance – the decision within levels	
			e.g At each level, as well as content, the scientific coherency of what is stated will help place the answer at the top, or the bottom, of that level.	
	0	No rewardable material.		
Level 1	1-2	Additional guidance	Possible candidate responses	
		Partially complete description of a suitable procedure with at least one measurement	Heat up the block in the boiling water. Then put the block into the cold water. Measure the temperature reached by the water.	
Level 2	3-4	Additional guidance	Possible candidate responses	
		Mostly complete description of a suitable procedure with at least two measurements and some description of processing the results.	As above with Measure mass of water. Use $\Delta Q = m \times c \times \Delta \theta$ to find thermal energy transferred	
Level 3	5-6	Additional guidance	Possible candidate responses	
suitab most measi clear		Detailed description of a suitable procedure with most of the necessary measurements and a clear description of processing the results.	As above with Calculate temperature changes by subtraction. Calculate thermal energy lost by Al as being equal to thermal energy gained by water.  Specific heat capacity of Al =	
			thermal energy transferred mass of Al × temp drop of Al	

#### Q11.

Question Number:	Answer	Additional guidance	Mark
	100 (°C) (1)	accept any answer between and including 95 and 102  (possibility that it is not pure water and possibility of heat loss prevents reaching boiling point)	(1) AO 2 1

#### Q12.

Question number	Answer	Additional guidance	Mark
	statements to include any <b>two</b> from		(2) AO1
	use cladding / (extra) insulation (1)		
	use double thicknesses of the concrete (1)	create cavity	
	use silver / reflective / white (paint) (1)		
	plant trees around (wind break) (1)		
	use double glazed windows (1)		
	(properly) close window(s)/door	draft exclusion	